



ACC.14

TCT@ACC-12 | innovation in intervention

A1569

JACC April 1, 2014

Volume 63, Issue 12



Stable Ischemic Heart Disease

COMPARATIVE SAFETY AND EFFICACY EVALUATION OF IVABRADINE, METOPROLOL AND ITS COMBINATION IN MANAGEMENT OF INAPPROPRIATE SINUS TACHYCARDIA IN CORONARY ARTERY BYPASS GRAFT PATIENTS

Poster Contributions

Hall C

Saturday, March 29, 2014, 3:45 p.m.-4:30 p.m.

Session Title: Old and Novel Therapies for Ischemic Heart Disease

Abstract Category: 26. Stable Ischemic Heart Disease: Therapy

Presentation Number: 1157-336

Authors: Niren Bhavsar, Roosha Parikh, Apurva Patel, Parth Parikh, Satya Gupta, Dhaval Naik, Chintan Mehta, Parloop Bhatt, Keyur Parikh, Dhiren Shah, Care Institute of Medical Science (CIMS), Ahmedabad, India, Cleveland Clinic, Cleveland, OH, USA

Background: Post-Operative Inappropriate Sinus Tachycardia (IST) is common After Coronary Artery Bypass Graft (CABG) surgery requiring treatment to reduce heart rate (HR). If untreated, increases incidence of complications including post-operative stroke.

Methods: A total of 151 patients undergoing CABG surgery at a tertiary care center in Western India from 10/2012 - 3/2013, who developed post-CABG IST, were divided as per cardiac surgeons discretion into 5 treatment groups for IST: Metoprolol 12.5 mg bid (n=47); Metoprolol 25 mg bid (n=24); Ivabradine 5 mg + Metoprolol 12.5 mg (bid,n=34); Ivabradine 5 mg + Metoprolol 25 mg (bid,n=23); and Ivabradine 5 mg bid (n=23). Safety and Efficacy were compared.

Results: Of the 151 patients (Mean age 59 ± 8 years, Male 87%), 41% had diabetes, 45% had hypertension and 67% had triple vessel disease. Mean pre-operative vitals were: HR 80, BP 125/78 mm Hg; respiratory rate 21/min and temperature 98.4 °F. "Off pump" CABG was done in 98% patients. During surgery, nor adrenaline was used in 87% patients, dobutamine in 2% patients and both in 11% patients. Mean intra-operative fluid balance was 988 ml. Average ICU and total hospital stay were 3.7 and 7.3 days respectively. IST developed post-CABG, averaging a 32% increase from baseline HR. HR was reduced similarly ($p > 0.05$) in all treatment groups.

Conclusions: Metoprolol, Ivabradine and its combination equally and effectively controlled IST in CABG patients without inducing hypotension or arrhythmia.




Figure: Heart Rate Response over Time.